



5 November 2018

ASX : ARV
FRANKFURT : ATY
US OTC : ARTTF

GOLD, COBALT AND COPPER IN THE WEST PILBARA

ARTEMIS RESOURCES LIMITED IS AN AUSTRALIAN MINERAL DEVELOPER ADVANCING ITS WEST PILBARA BASE, BATTERY AND PRECIOUS METALS ASSETS TOWARDS PRODUCTION.

ARTEMIS HAS CONSOLIDATED A MAJOR LAND HOLDING IN THE WEST PILBARA AND IS THE 100% OWNER OF THE RADIO HILL OPERATIONS AND PROCESSING INFRASTRUCTURE, STRATEGICALLY LOCATED 30 KM FROM THE CITY OF KARRATHA, THE POWERHOUSE OF THE PILBARA.

WANT TO KNOW MORE ABOUT ARTEMIS?

Please Contact:

Wayne Bramwell – Chief Executive Officer E. wayne.bramwell@artemisresources.com.au

E: Ed.Mead@artemisresources.com.au

David Tasker – Media Advisor
E: dtasker@chapteroneadvisors.com.au

Or visit the Artemis Website or follow us on Twitter.

Artemis Resources Limited

Suite 1, 11 Ventnor Ave, West Perth WA, Australia, 6005

P:+61863190000

E: info@artemisresources.com.au www.artemisresources.com.au Twitter - @Artemis_ARV

THREE NEW GOLD DISCOVERIES IN THE WEST PILBARA

11.4 g/t gold rock chip from Patterson's Hut (4.5km long anomaly)

Artemis Resources Limited ("Artemis" or "the Company") (ASX:ARV, Frankfurt: ATY, US OTC: ARTTF) is pleased to announce that the first regional scale compilation of geochemical data in this region has identified three (3) new discoveries within 30 km of Artemis's Radio Hill processing plant.

HIGHLIGHTS

GOLD - three new, large gold geochemical targets discovered:

- Patterson's Hut rock chip samples in veins and gossanous chert up to 11.4 g/t gold across a 4.5 km long surface soil anomaly
- Ruth Well North rock chip samples up to 5.04 g/t gold across a
 14 km long surface anomaly
- Pipeline soil geochemistry anomaly in association with numerous nuggets in shear zones across two 1.0 km long trends.

In addition, the Company has identified new extensions at the existing gold targets:

- Silica Hills new 1.5 km long gold in soil anomaly
- Nickol River strong gold geochemistry suggests additional primary targets may exist.

Artemis' Executive Director Ed Mead commented:

"Karratha until recently has remained relatively unexplored for its gold potential. This area wide geochemical and rock chip survey indicates substantial gold prospectivity exists within Artemis' extensive tenement package, all within a short trucking distance of our Radio Hill processing plant. The use of ionic geochemical analysis techniques in conjunction with traditional geochemical sampling has helped Artemis identify numerous new targets and projects such as Carlow Castle and Purdy's Reward."



REGIONAL GEOCHEMICAL PROGRAMME DEFINES 3 ENTIRELY NEW GOLD TARGETS

Artemis has undertaken a broad regional soil geochemistry sampling programme across the consolidated Artemis tenure on 400 metre spaced lines aligned north-south. Samples were taken 100m apart along these lines and a major multi-element suite of analyses were then conducted. A total of 12,247 samples have been collected.

Gold geochemistry responds strongly at Carlow Castle, Nickol River, Silica Hills, Pattersons Hut, Ruth Well North and Pipeline with the latter three areas being new discoveries.

All data presented in **Figure 1** below has been domained based on the GSWA 1:100,000 geological mapping, then ratio-ed using the 25th Percentile of the data. Data was contoured using Surfer software using Inverse distance squared (ID2) and the search ellipse long axis orientated to 80° east of north, contouring/plotting colours are then based on the 99th, 97.5th, 95th, 90th and 75th percentiles of the ratio-ed values.

The specific purpose of this processing was to highlight the anomalous samples and to minimise the lithological effects/contents of the differing underlying geological sequences.




Figure 1: Targets Identified or Highlighted by Regional Gold Geochemistry.






Figure 2: Soil Sampling Location and areas with Au related targets identified.

REGIONAL GOLD ENDOWMENT

Apart from the current high-profile conglomerate related gold occurrences at Purdy's Reward – Comet Well, the West Pilbara has a long history of small-scale gold production predominantly from quartz vein related systems. Artemis noted the presence of shear zone hosted gold at Nickol River feeding the alluvial/eluvial systems in the area and the axial plane quartz-gold-arsenopyrite mineralisation at the Weerianna Au Project.

After consideration of this information with the multiple other known gold sources within the greenstones, geochemical exploration was initiated in the Carlow Castle area.

Based on the Carlow Castle success, sampling was subsequently expanded to cover virtually all Artemis' tenure (Figure 2).



All the main areas identified in the geochemistry show multi-element responses as summarized below:

Carlow Castle - Au, Ag, Co, Cu, Ni, Hg, Mo, Se, Te, Pd, Zn

Patterson's Hut - Au, Ag, As, Mo, Ni, Sb, Se, Te, W

Ruth Well North - Au, Ag, Hg, Mo, Sb, W
Pipeline - Au, Ag, As, Co, Mo, Se, Tl

Silica Hills - Au, Ag, Bi, Mo, Sb Nickol River - Au, Ag, Hg, Mo, Se, Tl

Gold geochemistry responds strongly at Carlow Castle, Nickol River, Silica Hills, Pattersons Hut, Ruth Well North and Pipeline with the latter three areas previously unknown.

Patterson's Hut Au Prospect

Patterson's Hut is an entirely new area of gold in soil anomalism (to 68ppb Au) and mineralization. The soil geochemistry shows a continuous anomaly >95th percentile over **4.5km long** with an additional **1.7km** to the west after a small break (**Figure 1**).

Geological mapping (**Figure 3**) shows the area to be within a wide zone of sheared talcose and cherty schists with multiple strike parallel quartz veins, with rock chip sampling returning values up to 11.4 g/t Au. A discontinuous traverse of 20 rock chip samples over a width of 250m showed 9 samples with responses >1g/t Au to a maximum of 9.89 g/t Au.

Most samples were from quartz veins, but 3 samples were within gossanous chert, cherty gossan or chert with gossanous lenses which contained 1.41g/t, 1.32g/t and 2.41 g/t Au respectively (Appendix A). Interestingly many samples show a silver to gold ratio >10:1 possibly indicative of sulphide mineralisation.




Figure 3: Patterson's Hut Central Target with Gold in Soil and Rock Chip Values over 4.5km strike



Ruth Well North Au Prospect

Located to the north of the Ruth Well Ni-Cu mineralisation (**Figure 1**), a discontinuous gold in soil anomaly to a maximum 146ppb Au is traceable for **14 km**, apparently relating to a prominent chert ridge and outcrop. To date geological mapping and sampling has had limited success with one sample of a silicified laminated sediment containing 5.04g/t Au, the only sample of significance (**Appendix A**).

Historic Rotary Air blast (RAB) drilling 5.5km to the east of the main area along the siliceous laminated chert horizon (Figure 4) contained 1 sample of 0.5g/t Au over 1m.




Figure 4: Ruth Well North Central Target with Gold in Soil Values and Significant Rock Chips

Pipeline

Located to the south east of Radio Hill (**Figure 1**), soil geochemistry and metal detected nuggets with a Minelab GPZ 7000 defined 2 parallel trends approximately 1km apart, which are coincident with aeromagnetic trends (**Figure 5**). The aeromagnetic trends are interpreted to represent shear zones along the southern contact of the small Yannery Granite intrusion.

Outcrop in the area is subdued with metal detected nuggets being small angular fragments near quartz vein outcrops or scree, and are interpreted to represent the shear zones.






Figure 5: Pipeline Target with Gold in Soil Values and Nugget locations on RTPFVD Aeromagnetic Imagery

Silica Hills Au Prospect

This area is well known locally for producing coarse nuggety gold from quartz veins (**Figure 1**). The soil anomaly shows a zone **1.5km long** (>95th percentile) with a maximum value of 370ppb Au with probable extensions to the northeast and southwest. The anomalous element suite of Mo and Bi is suggestive of intrusive relationships.

Initial reverse circulation and rotary air blast drilling have been negative so the source of the anomalism is yet to be identified.

Nickol River

The gold geochemistry is surprisingly strong over the Nickol River alluvial area (Figure 1). This will be in part due to the extensive disturbance which has occurred in the area, but the geochemistry also suggests the known shear zones hosting primary mineralisation in the Samantha, Tozers and Boiler zones may be replicated elsewhere.

Historical drilling has been limited and only gold has been analysed from this drilling, hence making judgements about repetition or other mineralisation styles impossible until further assessment is undertaken.

Please refer to Appendix A for all significant rock chip sample results.





LOOKING FORWARD

Some infill sampling has been completed across the regional tenure and assays are pending. The Company is ranking and prioritising targets now with plans for first pass air-core drilling being developed following receipt of POW and heritage approvals.

For further information on this update or the Company generally, please visit our website at www.artemisresources.com.au or contact:

Edward Mead Executive Director Telephone: +61 407 445 351

Wayne Bramwell Chief Executive Officer Telephone: +61 417 953 073

David Tasker Advisor – Chapter One Telephone : +61 433 112 936





COMPETENT PERSONS STATEMENT

The information in this announcement that relates to Exploration Results and Exploration Targets is based on information compiled or reviewed by Allan Younger, who is a Member of the Australasian Institute of Mining and Metallurgy. Mr Younger is a consultant to the Company. Mr Younger has sufficient experience that is relevant to the style of mineralisation and type of deposit under consideration and to the activity which he is undertaking to qualify as a Competent Person as defined in the 2012 Edition of the 'Australasian Code for Reporting of Exploration Results, Mineral Resources and Ore Reserves'.

Mr Younger consents to the inclusion in the announcement of the matters based on his information in the form and context in which it appears.

BACKGROUND INFORMATION ON ARTEMIS RESOURCES

Artemis Resources Limited is an exploration and development company focussed on its large (≈2,400 km²) and prospective base, battery and precious metals assets in the Pilbara region of Western Australia. Artemis owns 100% of the 500,000 tpa Radio Hill processing plant and infrastructure, located approximately 35 km south of the city of Karratha.

The Company is evaluating 2004 and 2012 JORC Code compliant resources of gold, nickel, copper-cobalt, PGE's and zinc, all situated within a 40 km radius of the Radio Hill plant.

Artemis have signed Definitive Agreements with Novo Resources Corp. ("Novo"), which is listed on Canada's TSX Venture Exchange (TSXV:NVO), and pursuant to the Definitive Agreements, Novo has satisfied its expenditure commitment, and earned 50% of gold (and other minerals necessarily mined with gold) in conglomerate and/or paleoplacer style mineralization in Artemis' tenements within 100 km of the City of Karratha, including at Purdy's Reward ("the Gold Rights"). The Gold Rights do not include:

- (i) gold disclosed in Artemis' existing (at 18 May 2017) JORC Code Compliant Resources and Reserves; or
- (ii) gold which is not within conglomerate and/or paleoplacer style mineralization; or
- (iii) minerals other than gold.

Artemis' Mt Oscar tenement is excluded from the Definitive Agreements. The Definitive Agreements cover 36 tenements / tenement applications that are 100% owned by Artemis.

Pursuant to Novo's successful earn-in, two 50:50 joint ventures have been formed between Novo's subsidiary, Karratha Gold Pty Ltd ("Karratha Gold") and two subsidiaries of Artemis (KML No 2 Pty Ltd and Fox Radio Hill Pty Ltd). The joint ventures are managed as one by Karratha Gold with Artemis and Novo contributing to further exploration and any mining of the Gold Rights on a 50:50 basis.

FORWARD LOOKING STATEMENTS AND IMPORTANT NOTICE

This report contains forecasts, projections and forward-looking information. Although the Company believes that its expectations, estimates and forecast outcomes are based on reasonable assumptions it can give no assurance that these will be achieved. Expectations, estimates and projections and information provided by the Company are not a guarantee of future performance and involve unknown risks and uncertainties, many of which are out of Artemis' control.

Actual results and developments will almost certainly differ materially from those expressed or implied. Artemis has not audited or investigated the accuracy or completeness of the information, statements and opinions contained in this announcement. To the maximum extent permitted by applicable laws, Artemis makes no representation and can give no assurance, guarantee or warranty, express or implied, as to, and takes no responsibility and assumes no liability for the authenticity, validity, accuracy, suitability or completeness of, or any errors in or omission from, any information, statement or opinion contained in this report and without prejudice, to the generality of the foregoing, the achievement or accuracy of any forecasts, projections or other forward looking information contained or referred to in this report.

Investors should make and rely upon their own enquiries before deciding to acquire or deal in the Company's securities.



Appendix A: Rock Chip Assays

Sample ID	Easting	Northing	Description	Au ppm	Ag ppm	As ppm	Cu ppm	Ni ppm	Pb ppm
Patterson's Hut									
ARV025334	488027	7696026	QV; 0.5m thick, white, iron oxide stained	11.4	7.4	7	6	3	42
ARV025368	489121	7696666	QV; white, purple- red, grey, brecciated	9.89	139	2.5	38	2	1920
ARV025317	486407	7695762	QV; white, iron oxide stains, rare black blebs, old excavated hole in vein.	4.94	27.8	19	11	1	683
ARV025230	487021	7695845	QV; small 1m deep hole excavated in vein, white, bucky, 20m strike, 2m thick. No dip	4.59	65.8	9	27	7	1725
ARV025381	489085	7696827	QV; continuous from location 362, white, 50cm thick, sugary texture, minor brecciation	4.47	29.5	2.5	4	0.5	9
ARV025238	488515	7696530	Quartzite? and QV	4.38	15.9	16	291	9	153
ARV025301	486539	7696110	QV; 30cm thick striking 10m, banded, black- brown laminae	4.02	22.8	2.5	25	3	57
ARV025376	489123	7696735	QV blow; 2m thick, 10m radius, white, bucky, iron oxide stains	3.53	17.7	2.5	14	1	216
ARV025371	489143	7696624	QV; 30cm thick striking for 2m, brecciated, strong orange and purple iron oxide staining, sulfidic laminae	2.68	30.9	5	31	0.5	69
ARV025363	489118	7696703	QV; white- translucent, banded	2.47	29.4	2.5	28	1	112
ARV031913	489529	7696304	Chert; orange, banded, minor gossan lenses, strong iron oxide staining, brecciated in places	2.41	0.8	27	26	10	2
ARV025244	488705	7696693	QV; 50cm thick, strikes over 5m, white, brecciated, iron oxide staining, possible rare malachite.	2.23	6.2	2.5	183	3	354
ARV025377	489123	7696755	QV blow; 5m thick, 10m radius, orientated 002-182 with other QV blow outcrops.	1.83	46.1	5	41	0.5	1920



)

Sample ID	Easting	Northing	Description	Au ppm	Ag ppm	As ppm	Cu ppm	Ni ppm	Pb ppm
ARV025318	486554	7695744	QV; excavated holes in QV and host rock. Vein is white, bucky, rare black sulfidic blebs,	1.75	17.9	2.5	19	0.5	1335
ARV025335	488024	7695030	Altered QV host rock; cream-olive green, strongly silicified, brecciated, pervasive chlorite alteration. Possible association of QV with dolerite	1.43	0.25	7	7	10	14
ARV025261	490935	7696970	Gossanous chert; Taken along contact with green schist	1.41	1.3	1605	79	71	67
ARV025373	489098	7696649	Folded QV; south limb at 068, northern limb at 030, white, 50cm thick, bucky, minor brecciation	1.35	12.9	2.5	24	1	153
ARV025346	489568	7696544	Cherty Gossan; black-purple, vugs	1.32	0.25	906	41	198	24
ARV025370	489117	7696782	QV; 50cm thick striking for 10m, white-translucent, brecciated, iron oxide. Possibly folded	1.07	6	2.5	6	2	13
ARV025293	486880	7695819	QV; brecciated, purple-orange staining,	0.87	9.6	7	155	22	60
ARV025312	486590	7695829	QV; 40cm thick striking 3m, bucky with iron oxide stains.	0.67	3.3	8	4	19	3
ARV025284	489015	7696516	Phyllite; QV host to north	0.59	0.25	104	21	50	94
ARV025367	489123	7696670	QV blow; 3m thick, 8m strike, white, bucky, sulfidic veinlets and iron oxide/orange staining	0.56	11.1	6	9	2	305
ARV025242	488630	7696493	QV; up to 2m thick, striking over 5m, white, bucky with iron oxide laminations	0.45	1.1	7	5	5	82







Sample ID	Easting	Northing	Description	Au ppm	Ag ppm	As ppm	Cu ppm	Ni ppm	Pb ppm
Ruth Well									
ARV011559	485077	7693590	Laminated sediment. Quite siliceous. Some small cavities indicating possible sulfides.	5.04	50.9	735	261	61	39
ARV031784	485142	7693595	Ferruginous chert outcrop. Sample has slight gossanous appearance.	0.54	1.5	659	60	291	23
ARV031760	485404	7693458	Siliceous. More like a flint than a chert. Big boulders, not in situ.	0.47	1.2	75	40	18	10



JORC Code, 2012 Edition - Table 1

Section 1 Sampling Techniques and Data

(Criteria in this section apply to all succeeding sections.)

Criteria	JORC Code explanation	Commentary
Sampling techniques	 Nature and quality of sampling (e.g. cut channels, random chips, or specific specialised industry standard measurement tools appropriate to the minerals under investigation, such as down hole gamma sondes, or handheld XRF instruments, etc). These examples should not be taken as limiting the broad meaning of sampling. Include reference to measures taken to ensure sample representivity and the appropriate calibration of any measurement tools or systems used. Aspects of the determination of mineralisation that are Material to the Public Report. In cases where 'industry standard' work has been done this would be relatively simple (e.g. 'reverse circulation drilling was used to obtain 1 m samples from which 3 kg was pulverised to produce a 30 g charge for fire assay'). In other cases, more explanation may be required, such as where there is coarse gold that has inherent sampling problems. Unusual commodities or mineralisation types (e.g. submarine nodules) may warrant disclosure of detailed information. 	 The soil samples were uniformly collected from 15cm, with colour, moisture and general topography recorded. Two forms of analysis were conducted for the soils conventional analysis using the AuME-ST44 was applied to samples sieved to -2mm. The second method was Ionic leach where soil samples are sieved to -4mm. The AuME-ST44 is an aqua regia digest with ICP-MS finish for multi-element analysis including: Au, Ag, Al, As, Ba, Be Bi, Ca, Cd, Co, Cr, Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P Pb, Pd, Pt, S, Sb, Sc, Sr, Th, Ti, Tl, U, V, W, Zn. Samples are pulverised to 95% passing 75 microns for maximum digestion. Ionic Leach™ uses a cyanide leach in a buffered solution digest with ICP-MS finish for ultra-low level detection levels for elements including: Au, Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co Cr, Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P, Pb, Pd, Pt, S, Sb Sc, Sr, Th, Ti, Tl, U, V, W, Zn, REE. Samples are unpulverized and the technique is known as a partial extraction approach. Field duplicates were taken and submitted for analysis with the soil samples. Rock chip samples were pulverised to 95% passing 75 microns, Au by 50-gram Fire Assay (Au-AA26) with ICF finish. Multi element analysis used 4 Acid Digest ICP-AES Finish (ME-ICP61) for: Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co, Cr Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P, Pb, S, Sb, Sc, Sr, Th Ti, Tl, U, V, W, Zn. Rock chip samples from the Bel's PGE area were analysed using the AuME-ST44 technique
Drilling techniques	 Drill type (e.g. core, reverse circulation, open-hole hammer, rotary air blast, auger, Bangka, sonic, etc) and details (e.g. core diameter, triple or standard tube, depth of diamond tails, face-sampling bit or other type, whether core is oriented and if so, by what method, etc). 	Not drilling data
Drill sample recovery	 Method of recording and assessing core and chip sample recoveries and results assessed. Measures taken to maximise sample recovery and ensure representative nature of the samples. Whether a relationship exists between sample recovery and grade and whether sample bias may have occurred due to preferential loss/gain of fine/coarse material. 	Not drilling data.
Logging	 Whether core and chip samples have been geologically and geotechnically logged to a level of detail to support appropriate Mineral Resource estimation, mining studies and metallurgical 	Rock chip samples were logged.



RTEMIS		ASX / MEDIA ANNOUNCEMENT
Criteria	JORC Code explanation	Commentary
	 Whether logging is qualitative or quantitative in nature. Core (or costean, channel, etc) photography. The total length and percentage of the relevant intersections logged. 	
Sub-sampling techniques and sample preparation	 If core, whether cut or sawn and whether quarter, half or all core taken. If non-core, whether riffled, tube sampled, rotary split, etc and whether sampled wet or dry. For all sample types, the nature, quality and appropriateness of the sample preparation technique. Quality control procedures adopted for all subsampling stages to maximise representivity of samples. Measures taken to ensure that the sampling is representative of the in-situ material collected, including for instance results for field duplicate/second-half sampling. Whether sample sizes are appropriate to the grain size of the material being sampled. 	 Not drilling data Duplicate samples were collected and submitted for analysis with Reference standards inserted during soil sampling. Reference samples were inserted with rock chip samples.
Quality of assay data and laboratory tests	 The nature, quality and appropriateness of the assaying and laboratory procedures used and whether the technique is considered partial or total. For geophysical tools, spectrometers, handheld XRF instruments, etc, the parameters used in determining the analysis including instrument make and model, reading times, calibrations factors applied and their derivation, etc. Nature of quality control procedures adopted (e.g. standards, blanks, duplicates, external laboratory checks) and whether acceptable levels of accuracy (i.e. lack of bias) and precision have been established. 	 ALS (Perth) were used for all analysis of samples submitted by Artemis. The laboratory techniques below are for all samples submitted to ALS and are considered appropriate for the styles of mineralisation within the Karratha region: The AuME-ST44 is an aqua regia digest with ICP-MS finish for multi-element analysis including: Au, Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co, Cr, Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P, Pb, Pd, Pt, S, Sb, Sc, Sr, Th, Ti, Tl, U, V, W, Zn. Samples are pulverised to 95% passing 75 microns for maximum digestion. Ionic Leach™ uses a cyanide leach in a buffered solution digest with ICP-MS finish for ultra-low level detection levels for elements including: Au, Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co, Cr, Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P, Pb, Pd, Pt, S, Sb, Sc, Sr, Th, Ti, Tl, U, V, W, Zn, REE. Samples are unpulverized and the technique is known as a partial extraction approach. Field duplicates were taken and submitted for analysis with the soil samples. Rock chip samples were pulverised to 95% passing 75 microns for maximum digestion, Au by 50-gram Fire Assay (Au-AA26) with ICP finish. Multi element analysis used 4 Acid Digest ICP-AES Finish (ME-ICP61) for: Ag, Al, As, Ba, Be, Bi, Ca, Cd, Co, Cr, Cu, Fe, Ga, K, La, Mg, Mn, Mo, Na, Ni, P, Pb, S, Sb, Sc, Sr, Th, Ti, Tl, U, V, W, Zn. Rock chip samples from the Bel's PGE area were analysed

using the AuME-ST44 technique

Artemis.

Artemis.

Standards were used for external laboratory checks by

Duplicates were used for external laboratory checks by



ASX / MEDIA ANNOUNCEMENT

RESOURCES		
Criteria	JORC Code explanation	Commentary
Verification of sampling and assaying	 The verification of significant intersections by either independent or alternative company personnel. The use of twinned holes. Documentation of primary data, data entry procedures, data verification, data storage (physical and electronic) protocols. Discuss any adjustment to assay data. 	At least two company personnel verify all significant results.
Location of data points	 Accuracy and quality of surveys used to locate drill holes (collar and down-hole surveys), trenches, mine workings and other locations used in Mineral Resource estimation. Specification of the grid system used. Quality and adequacy of topographic control. 	 location of the soil and rock chip samples. The grid system used for all Artemis sampling is GDA94 (MGA 94 Zone 50)
Data spacing and distribution	 Data spacing for reporting of Exploration Results. Whether the data spacing and distribution is sufficient to establish the degree of geological and grade continuity appropriate for the Mineral Resource and Ore Reserve estimation procedure(s) and classifications applied. Whether sample compositing has been applied. 	Not drill data.
Orientation of data in relation to geological structure	 Whether the orientation of sampling achieves unbiased sampling of possible structures and the extent to which this is known, considering the deposit type. If the relationship between the drilling orientation and the orientation of key mineralised structures is considered to have introduced a sampling bias, this should be assessed and reported if material. 	 Not drill data. Geochemical sampling has been undertaken on a nominal 400m x 100m spacing.
Sample security	The measures taken to ensure sample security.	 The chain of custody is managed by the supervising geologist who places calico sample bags in polyweave sacks. Up to 10 calico sample bags are placed in each sack. Each sack is clearly labelled with: Artemis Resources Ltd Address of laboratory Sample range Samples were delivered by Artemis personnel to the transport company in Karratha and shrink wrapped onto pallets. The transport company then delivers the samples directly to the laboratory.
Audits or reviews	The results of any audits or reviews of sampling techniques and data.	Data is validated upon up-loading into the master database. Any validation issues identified are investigated prior to reporting of results.



Section 2 Reporting of Exploration Results

(Criteria listed in the preceding section also apply to this section.)

Criteria	JORC Code explanation	Commentary
Mineral tenement and land tenure status	 Type, reference name/number, location and ownership including agreements or material issues with third parties such as joint ventures, partnerships, overriding royalties, native title interests, historical sites, wilderness or national park and environmental settings. The security of the tenure held at the time of reporting along with any known impediments to obtaining a licence to operate in the area. 	The regional soil sampling has been conducted across virtually all Artemis and subsidiary's tenements.
Exploration done by other parties	 Acknowledgment and appraisal of exploration by other parties. 	 Previous explorers in the region include but not limited to are Westfield Minerals, Consolidated Gold Areas, Open Pit Mining and Exploration, Legend Mining, Agip Exploration, Titan Resources and Fox Resources.
Geology	 Deposit type, geological setting and style of mineralisation. 	The soil sampling program was planned identifying any unknown styles of mineralization in the West Pilbara.
Drill hole Information	 A summary of all information material to the understanding of the exploration results including a tabulation of the following information for all Material drill holes: easting and northing of the drill hole collar elevation or RL (Reduced Level – elevation above sea level in metres) of the drill hole collar dip and azimuth of the hole down hole length and interception depth hole length. If the exclusion of this information is justified on the basis that the information is not Material and this exclusion does not detract from the understanding of the report, the Competent Person should clearly explain why this is the case. 	Not drilling data.
Data aggregation methods	 In reporting Exploration Results, weighting averaging techniques, maximum and/or minimum grade truncations (e.g. cutting of high grades) and cut-off grades are usually Material and should be stated. Where aggregate intercepts incorporate short lengths of high-grade results and longer lengths of low-grade results, the procedure used for such aggregation should be stated and some typical examples of such aggregations should be shown in detail. The assumptions used for any reporting of metal equivalent values should be clearly stated. 	Not drilling data.
Relationship between mineralisation	 These relationships are particularly important in the reporting of Exploration Results. If the geometry of the mineralisation with respect to the drill hole angle is known, its 	Not drilling data



ASX / MEDIA ANNOUNCEMENT

RESOURCES		
Criteria	JORC Code explanation	Commentary
widths and	nature should be reported.	
intercept lengths	 If it is not known and only the down hole lengths are reported, there should be a clear statement to this effect (e.g. 'down hole length, true width not known'). 	
Diagrams	 Appropriate maps and sections (with scales) and tabulations of intercepts should be included for any significant discovery being reported These should include, but not be limited to a plan view of drill hole collar locations and appropriate sectional views. 	 Appropriate maps and sections are available in the body of this announcement.
Balanced reporting	 Where comprehensive reporting of all Exploration Results is not practicable, representative reporting of both low and high grades and/or widths should be practiced to avoid misleading reporting of Exploration Results. 	Reporting of results in this report is considered balanced.
Other substantive exploration data	Other exploration data, if meaningful and material, should be reported including (but not limited to): geological observations; geophysical survey results; geochemical survey results; bulk samples – size and method of treatment; metallurgical test results; bulk density, groundwater, geotechnical and rock characteristics; potential deleterious or contaminating substances.	The regional soil exploration program was to establish the baseline information.
Further work	 The nature and scale of planned further work (e.g. tests for lateral extensions, depth extensions or large-scale step-out drilling). Diagrams clearly highlighting the areas of possible extensions, including the main geological interpretations and future drilling areas, provided this information is not commercially sensitive. 	Follow-up of the numerous identified anomalous areas will continue.